
Androvite® is intended to provide proper nutrition for adult men.\* It is a magnesium-emphasized nutritional supplement that may be used as part of a total dietary program, which consists of change in lifestyle, dietary guidelines, exercise, and Androvite®.

**SUGGESTED USE:** As a dietary supplement, 6 tablets daily with meals, or as directed by a healthcare professional.

Serving Size	6 tablets
Servings per container	30
Amount per serving:	
Vitamin A (as retinyl palmitate and 80% as beta carotene)	25,000 IU
Vitamin C (as ascorbic acid)	1000 mg
Vitamin D (as cholecalciferol)	400 IU
Vitamin E (as d-alpha tocopheryl succinate)	400 IU
Vitamin B1 (as thiamine mononitrate)	50 mg
Vitamin B2, Riboavin	50 mg
Niacin (as niacinimide)	50 mg
Vitamin B6 (as pyridoxine HCI)	100 mg
Folic Acid	400 μg
Vitamin B12 (as hydroxyocobalamin)	125 μg
Biotin	125 μg
Pantothenic Acid (as calcium pantothenate)	100 mg
Iron (as amino acid chelate)	18 mg
Iodine (as hydrolyzed protein complex)	150 μg
Magnesium (as magnesium oxide)	500 mg
Zinc (as amino acid chelate)	50 mg
Selenium (as hydrolyzed protein complex)	200 μg
Copper (as amino acid chelate)	2 mg
Manganese (as amino acid chelate)	10 mg
Chromium (as hydrolyzed protein complex)	200 μg
p-Aminobenzoic Acid (PABA)	25 mg
Betaine HCI	100 mg
Pancreatin 4X	75 mg
Inositol	36 mg
Hesperidin	35 mg
Rutin	25 mg
Boron (hydrolyzed protein complex)	3 mg



**Other ingredients:** Micosolle®, a silica-based excipient containing a non-ionic surfactant, cellulose and modied cellulose, silica, stearic acid, pharmaceutical glaze, magnesium stearate, natural flavor and annatto as natural source of color.

Taking more than 400  $\mu$ g of selenium per day from all sources should only be done under the guidance of a healthcare professional. Do not exceed 6 tablets a day. Do not use for pernicious anemia, or if you have hemochromatosis or hemosiderosis.

**WARNING:** Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or poison control center immediately.